

Propositional Logic: Semantics (3/3)

CS402, Spring 2018

Shin Yoo

- Semantic Tableaux
- Soundness and completeness

A relatively efficient algorithm for deciding satisfiability in the propositional calculus.

- Search systematically for a model.
- If one is found, the formula is satisfiable; otherwise, it is unsatisfiable.

This method is the main tool for proving general theorems about the calculus.

Definition 1 (2.57)

A *literal* is an atom or a negation of an atom. An atom is a positive literal and the negation of an atom is a negative literal. For any atom p , $\{p, \neg p\}$ is a *complementary* pair of literals. For any formula A , $\{A, \neg A\}$ is a *complementary* pair of formulas. A is the complement of $\neg A$ and $\neg A$ is the complement of A .

Important observation: a set of literals is *satisfiable* if and only if it does **not** contain a *complementary* pair of literals.

Semantic Tableaux

Analyze the satisfiability of $A = p \wedge (\neg q \vee \neg p)$ in an arbitrary interpretation \mathcal{I} .

$$\nu_{\mathcal{I}}(A) = T \text{ iff both } \nu_{\mathcal{I}}(p) = T \text{ and } \nu_{\mathcal{I}}(\neg q \vee \neg p) = T.$$

Hence, $\nu_{\mathcal{I}}(A) = T$ if and only if either:

- ① $\nu_{\mathcal{I}}(p) = T$ and $\nu_{\mathcal{I}}(\neg q) = T$ or
- ② $\nu_{\mathcal{I}}(p) = T$ and $\nu_{\mathcal{I}}(\neg p) = T$

$\therefore A$ is satisfiable if and only if there exists an interpretation such that (1) holds or (2) holds.

The process is to reduce the question from one about the satisfiability of a formula to one about the satisfiability of sets of *literals*. Since any formula contains *finite* atoms, there are at most *finite* number of sets of literals. Then the decision on satisfiability becomes trivial.

Formula $B = (p \vee q) \wedge (\neg p \wedge \neg q)$ under an arbitrary interpretation \mathcal{I} .

$\nu_{\mathcal{I}}(B) = T$ iff $\nu_{\mathcal{I}}(p \vee q) = T$ and $\nu_{\mathcal{I}}(\neg p \wedge \neg q) = T$.

Hence, $\nu_{\mathcal{I}}(B) = T$ iff $\nu_{\mathcal{I}}(p \vee q) = \nu_{\mathcal{I}}(\neg p) = \nu_{\mathcal{I}}(\neg q) = T$.

Hence, $\nu_{\mathcal{I}}(B) = T$ iff either:


- 1 $\nu_{\mathcal{I}}(p) = \nu_{\mathcal{I}}(\neg p) = \nu_{\mathcal{I}}(\neg q) = T$, or
- 2 $\nu_{\mathcal{I}}(q) = \nu_{\mathcal{I}}(\neg p) = \nu_{\mathcal{I}}(\neg q) = T$.

Since both $\{p, \neg p, \neg q\}$ and $\{q, \neg p, \neg q\}$ contain complementary pairs, B is unsatisfiable.


- This systematic search becomes easier if we use a suitable data structure to keep track of the assignments that must be made to subformulas.
- In semantic tableaux, trees are used.
- A leaf containing a complementary set of literals will be marked with a \times symbol, while a leaf containing a satisfiable set of literals will be marked with a \odot symbol.

Semantic tableaux



Is $p \wedge (\neg q \vee \neg p)$ satisfiable?



Is $(p \vee q) \wedge (\neg p \wedge \neg q)$ satisfiable?



The tableau construction is not unique.



Semantic tableaux

Classification of formulas according to their **principal operators**:

- α -formulas are conjunctive and are satisfiable only if both subformulas, α_1 and α_2 , are satisfied.
- β -formulas are disjunctive and are satisfied if at least one of the subformulas, β_1 or β_2 , is satisfiable.

α	α_1	α_2
$\neg\neg A_1$	A_1	
$A_1 \wedge A_2$	A_1	A_2
$\neg(A_1 \vee A_2)$	$\neg A_1$	$\neg A_2$
$\neg(A_1 \rightarrow A_2)$	A_1	$\neg A_2$
$\neg(A_1 \uparrow A_2)$	A_1	A_2
$A_1 \downarrow A_2$	$\neg A_1$	$\neg A_2$
$A_1 \leftrightarrow A_2$	$A_1 \rightarrow A_2$	$A_2 \rightarrow A_1$
$\neg(A_1 \oplus A_2)$	$A_1 \rightarrow A_2$	$A_2 \rightarrow A_1$

β	β_1	β_2
$\neg(B_1 \wedge B_2)$	$\neg B_1$	$\neg B_2$
$B_1 \vee B_2$	B_1	B_2
$B_1 \rightarrow B_2$	$\neg B_1$	B_2
$B_1 \uparrow B_2$	$\neg B_1$	$\neg B_2$
$\neg(B_1 \downarrow B_2)$	B_1	B_2
$\neg(B_1 \leftrightarrow B_2)$	$\neg(B_1 \rightarrow B_2)$	$\neg(B_2 \rightarrow B_1)$
$B_1 \oplus B_2$	$\neg(B_1 \rightarrow B_2)$	$\neg(B_2 \rightarrow B_1)$

Let \mathcal{T} for a propositional formula A be a tree, whose nodes are all labeled with a set of formulas. Let $U(l)$ be the set of formulas of leaf l .

CONSTRUCTION OF SEM. TAB. (Algorithm 2.64)

Input: A propositional formula A

Output: A semantic tableaux \mathcal{T} for A with marked leaves

- (1) $\mathcal{T} \leftarrow$ a tree with a single node labeled $\{A\}$
- (2) **while** there exists an unmarked leaf
- (3) **foreach** unmarked leaf l
- (4) **if** $U(l)$ is a set of lit.
- (5) **if** a compl. lit. pair $\in U(l)$ **then** Mark l as \times
- (6) **else** Mark l as \oplus
- (7) **else**
- (8) Choose $A \in U(l)$
- (9) **if** $A \equiv \alpha$ **then** Add l' to l s.t. $U(l') \leftarrow (U(l) - \{\alpha\}) \cup \{\alpha_1, \alpha_2\}$
- (10) **if** $A \equiv \beta$ **then** Add l', l'' to l s.t. $U(l') \leftarrow (U(l) - \{\beta\}) \cup \{\beta_1\}$,
 $U(l'') \leftarrow (U(l) - \{\beta\}) \cup \{\beta_2\}$

This is not deterministic due to the choice of leaves in line (3).

Definition 2 (2.65)

- A tableau whose construction has terminated is called a *completed tableau*.
- A completed tableau is *closed* if all leaves are marked closed (i.e. \times); otherwise, it is *open*.

Theorem 1 (2.66)

The construction of a semantic tableau terminates.

Soundness and Completeness

A tool operates on a formula ϕ at the syntactic level, i.e. it does not apply all possible interpretations.

- A tool is *sound* if whenever the tool says that a formula ϕ is valid (validity, not satisfiability), ϕ is really valid. That is, $\vdash \phi$ implies $\models \phi$.
- A tool is *complete* if whenever ϕ is valid, the tool does say that ϕ is valid. That is, $\models \phi$ implies $\vdash \phi$.
 - Writing in a contra-positive way: a tool (or method) is complete if whenever the tool says that ϕ is not valid, then ϕ is really not valid.
- Therefore, if a tool is sound and complete, then the tool says that ϕ is valid iff ϕ is really valid.

Note that:

- If a dumb tool always says that ϕ is not valid, then that tool is still sound.
- If a dumb tool always says that ϕ is valid, then that tool is still complete.

Soundness and Completeness

Theorem 2 (2.67)

Let \mathcal{T} be a completed tableau for a formula A . A is unsatisfiable if and only if \mathcal{T} is closed.

Corollary 1 (2.68)

A is satisfiable if and only if \mathcal{T} is open.

Corollary 2 (2.69)

A is valid if and only if the tableau for $\neg A$ is closed.

Corollary 3 (2.70)

The method of semantic tableaux is a decision procedure for validity in the propositional calculus.

Proof of soundness:

- If the tableau \mathcal{T} for a formula A closes, then A is unsatisfiable.
- If a subtree rooted at node n of \mathcal{T} closes, then the set of formulas $U(n)$ labeling n is unsatisfiable. Let h be the height of the node n in \mathcal{T} .
 - If $h = 0$, n is a leaf. Since \mathcal{T} closes, $U(n)$ contains a complementary set of literals. Hence $U(n)$ is unsatisfiable.

- If $h > 0$, either α - or β - rule was used in creating the child(ren) of n :
 - Case 1: α -rule. $U(n) = \{A_1 \wedge A_2\} \cup U_0$ and $U(n') = \{A_1, A_2\} \cup U_0$ for some set of formulas U_0 .
 - The height of n' is $h - 1$; by induction, $U(n')$ is unsatisfiable since the subtree rooted at n' closes.
 - Let ν be an arbitrary interpretation. Since $U(n')$ is unsatisfiable, $\nu(A') = F$ for some $A' \in U(n')$. There are three possibilities:
 - For some $A_0 \in U_0$, $\nu(A_0) = F$. But $A_0 \in U_0 \subseteq U(n)$.
 - $\nu(A_1) = F$, $\nu(A_1 \wedge A_2) = F$. And $A_1 \wedge A_2 \in U(n)$.
 - $\nu(A_2) = F$, $\nu(A_1 \wedge A_2) = F$. And $A_1 \wedge A_2 \in U(n)$.

In all three cases, $\nu(A) = F$ for some $A \in U(n)$. Therefore, $U(n)$ is unsatisfiable.

- If $h > 0$, either α - or β - rule was used in creating the child(ren) of n :
 - Case 2: β -rule. $U(n) = \{B_1 \vee B_2\} \cup U_0$, $U(n) = \{B_1\} \cup U_0$ and $U(n'') = \{B_2\} \cup U_0$ for some set of formulas U_0 .
 - By induction, both $U(n')$ and $U(n'')$ are unsatisfiable, since the subtrees rooted at n' and n'' close.
 - Let ν be an arbitrary interpretation. There are three possibilities:
 - $U(n')$ and $U(n'')$ are unsatisfiable, because $\nu(B_0) = F$ for some $B_0 \in U_0$. But $B_0 \in U_0 \subseteq U(n)$.
 - Otherwise, $\nu(B_0) = T$ for all $B_0 \in U_0$. Since both $U(n')$ and $U(n'')$ are unsatisfiable, $\nu(B_1) = \nu(B_2) = F$. By definition of ν on \vee , $\nu(B_1 \vee B_2) = F$, and $B_1 \vee B_2 \in U(n)$.

Therefore $\nu(B) = F$ for some $B \in U(n)$; since ν was arbitrary, $U(n)$ is unsatisfiable.

Proof of completeness:

- If A is unsatisfiable, then every tableau for A closes.
- Contrapositive statement (Cor 2.68): if some tableau for A is open (i.e., if some tableau for A has an open branch), then the formula A is satisfiable.

Definition 3 (2.75)

Let U be a set of formulas. U is a **Hintikka**^a set iff:

- 1 For all atoms p appearing in a formula of U , either $p \notin U$ or $\neg p \notin U$.
- 2 If $\alpha \in U$ is an α -formula, then $\alpha_1 \in U$ and $\alpha_2 \in U$.
- 3 If $\beta \in U$ is an β -formula, then either $\beta_1 \in U$ or $\beta_2 \in U$.

^aNamed after Finnish logician Jaakko Hintikka (1929-2015).

Let us first deal with the following theorem, which we will then use to prove the completeness.

Theorem 3 (2.77)

Let l be an open leaf in a completed tableau \mathcal{T} . Let $U = \bigcup_i U(i)$, where i runs over the set of nodes on the branch from the root to l . Then U is a Hintikka set.

Proof.

Literal p or $\neg p$ cannot be decomposed. Thus, if a literal p or $\neg p$ appears for the first time in $U(n)$ for some n , the literal will be copied into $U(k)$ for all nodes k on the branch from n to l , in particular, $p \in U(l)$ or $\neg p \in U(l)$. This means that all literals in U appear in $U(l)$. Since the branch is open, no complementary pair of literals appears in $U(l)$, so Condition (1) for Hintikka set holds. \square

Proof for Theorem 2.77 Cont.

Suppose that $A \in U$ is an α -formula. Since the tableau is completed, A was the formula selected for decomposing at some node n in the branch from the root to l . Then $\{A_1, A_2\} \subseteq U(n') \subseteq U$, so Condition (2) holds.

Suppose that $B \in U$ is an β -formula. Since the tableau is completed, B was the formula selected for decomposing at some node n in the branch from the root to l . Then either $B_1 \in U(n') \subseteq U$ or $B_2 \in U(n') \subseteq U$, so Condition (3) holds. \square

Theorem 4 (2.78)

Hintikka's Lemma: *Let U be a Hintikka set. Then U is satisfiable.*

Proof.

Let us define an interpretation \mathcal{I} based on the fact that U is a Hintikka set, and then show \mathcal{I} is a model of U .

Let $\mathcal{I} : \mathcal{P}_U \rightarrow \{T, F\}$ be:

- $\mathcal{I}(p) = T$ if $p \in U$
- $\mathcal{I}(p) = F$ if $\neg p \in U$
- $\mathcal{I}(p) = T$ if $p \notin U$ and $\neg p \notin U$

Condition (1) in Definition 2.75 states that every literal is given exactly one value. The third case assigns arbitrary T to atoms that appear in U but not in literal form (i.e. $q \in \mathcal{P}_U$ but $q \notin U$ and $\neg q \notin U$). □

Proof for Theorem 2.78 Cont.

We use structural induction to show that for any $a \in U$, $\nu_{\mathcal{I}}(A) = T$. The base case is when A is an atom.

- A is an atom: $\nu_{\mathcal{I}}(A) = \nu_{\mathcal{I}}(p) = \mathcal{I}(p) = T$, because $p \in U$.
- A is a negated atom $\neg p$: $\neg p \in U$, therefore $\mathcal{I}(p) = F$, therefore $\nu_{\mathcal{I}}(A) = \nu_{\mathcal{I}}(\neg p) = T$.
- A is an α -formula: by Condition (2) of Def. 2.75, $A_1 \in U$ and $A_2 \in U$. By inductive hypothesis, $\nu_{\mathcal{I}}(A_1) = \nu_{\mathcal{I}}(A_2) = T$, so by definition of the conjunctive operator, $\nu_{\mathcal{I}}(A) = T$.
- A is an β -formula: by Condition (3) of Def. 2.75, either $B_1 \in U$ or $B_2 \in U$. By inductive hypothesis, either $\nu_{\mathcal{I}}(B_1) = T$ or $\nu_{\mathcal{I}}(B_2) = T$, so by definition of the disjunctive operator, $\nu_{\mathcal{I}}(A) = \nu_{\mathcal{I}}(B) = T$.



Proof of Completeness for Semantic Tableaux.

Let \mathcal{T} be a completed *open* tableau for A . Then U , the union of the labels of the nodes on *an open branch*, is a Hintikka set by Theorem 2.77, and a model can be found for U by Theorem 2.78. Since A is the formula labeling the root, $A \in U$, so the interpretation is a model of A . □